

Lingual Thyroid-Literature Review

Clara Rafael Silva Xavier¹, Lucas de Faria Barros Medeiros¹, Artur Dantas Freire¹, Thiago Emanuel Vêras Lemos¹, Amália Cínthia Meneses Rêgo² and Irami Araújo-Filho^{3*}

¹Graduating Medical University Potiguar – Laureate International Universities, Natal, Brazil

²Health University Potiguar - Laureate International Universities, Natal, Brazil

³Department of Surgery, University Potiguar - Laureate International Universities, Rio Grande do Norte Federal University, Natal, Brazil

*Corresponding author: Irami Araújo-Filho, Universidade Potiguar Natal, Rio Grande Do Norte Brazil, Tel: (84) 98876-0206; E-mail: irami.filho@uol.com.br

Rec date: January 22, 2016; Acc date: February 18, 2016; Pub date: February 29, 2016

Copyright: © 2016 Xavier CRS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The presence of ectopic thyroid tissue is a rare entity. The non-gland migration occurs in the early stages of embryogenesis to your normal cervical location. Thus, the ectopic tissue lodges in general in the path of the thyroglossal duct, in the middle line of the neck. The most common location is in the lingual zone, being called a lingual thyroid. This, in most cases will be asymptomatic. However, it is able to manifest itself with symptoms of dysphagia, dysphonia, obstruction of upper airways or hemorrhage at any moment between childhood and adulthood. This article is a review of this disease, targeting mainly the conduct, still very controversial in world literature.

Keywords: Ectopic thyroid; Lingual thyroid; Lingual goiter; Robotic surgery; Transoral surgery

Introduction

The ectopic thyroid was first described by Hickman in 1869, being defined as any thyroid tissue not found in their usual topography. The clinical incidence varies between 1:3000 and 1:100000, constituting in 70 to 100% of the occurrences, the only thyroid tissue is available. More common in women, with an incidence of 4:1:00 pm for men. Is the main cause of hypothyroidism in children, accounting for 52% of dysgenesis thyroid disease [1,2].

The presence of ectopic thyroid gland at the base of the tongue, referred to as lingual thyroid, is a rare congenital anomaly resulting from the incomplete descent of the gland during embryogenesis. Microscopic quantities are found in up to 10% of the population, however only 0.01% of these patients will have thyroid volume enough to make them symptomatic. As well as the thyroid normal location, the ectopic tissue will be subject to any thyroid pathology such as goiter or neoplasia [3-5].

90% cases of ectopic are lingual thyroid. Displays two peaks, to age 12 and 50 years of age. However, the age range of occurrence is quite broad, ranging from 6 to 74 years. 70 the 80% of patients do not have functional thyroid tissue elsewhere, plus 33% of them being with hypothyroidism. However, most will be euthyroid and a few euthyroid with hyperthyroidism [6-8].

The main differential diagnoses are: hemangioma of tongue, thyroglossal duct cyst, dermoid cyst, lipoma, lymphadenopathy, lymphangioma, cystic hygroma, besides neoplasms. However, the malignant transformation is rare and is usually diagnosed after surgical excision of the lesion, the Histopathological examination, through being the most papillary carcinomas or mixed carcinomas [2,3].

The lingual thyroid, although rare, clinically should be a relevant diagnostic hypothesis in the differential diagnosis of tumor structure medians of the base of the tongue. Seems to be more prevalent in the general population than the rare paintings reported to believe, being misdiagnosed, since it expresses, in necropsy studies, an incidence of 10%. By having a large number of therapeutic options available and few studies in the area, its treatment still could not be consolidated. So, this review aims to throw light on this little discussed in Medical Academy, in addition to promoting an update about the new treatments for this disease [4,5].

Methods

Consists of a literature review in the databases PubMed, MEDLINE, SCIELO, LILACS and SCOPUS, using lingual thyroid descriptors, robotic surgery, ectopic thyroid and lingual goiter. We included English and Portuguese language articles published between 1990 and 2015.

Pathophysiology

Thyroid embryonic development occurs between the third and the seventh week of pregnancy, originating on the floor of the pharynx of primitive and migrating earlier and inferiorly until it reaches its final location in the medial and anterior

neck, just below the cricoid cartilage. The ectopic thyroid is triggered by failure to complete or incomplete tissue migration during this period, and can be located at any point between the blind foramen and caudal part of the neck. Consequently, the most common locations are in lingual region, sublingual, prelaryngeal and substernal. Accommodation at other sites outside this path were also reported, as in superior mediastinum, pericardial sac, heart and malik [3,9].

The most common presentation between the thyroid is lingual ectopic, which usually is situated between the circumvallate papillae and the epiglottis, blind foramen. Its pathogenesis is not completely understood, however, it is speculated that maternal antibodies anti-thyroid hamper the descent of thyroid tissue during embryogenesis. Hypothyroidism could be justified by the same mechanism. Genetic changes and family history were also correlated to the origin of the frame [3,4].

Lingual thyroid is not commonly associated with anatomically at parathyroid, because the latter have an embryonic origin distinct from the dorsal wings of third and fourth pharyngeal pouches [4,7].

Clinical Presentation and Diagnosis

The lingual thyroid should always be investigated to find if tumor structure at the base of tongue, for it is the benign mass more often found in the Union of 2/3 1/3 previous back of tongue. Because it is often asymptomatic, is usually diagnosed by chance or in the study of clinical hypothyroidism. Will be symptomatic usually in situations where there is hypertrophy of the gland by increasing the metabolic needs, which will be stimulated by the excess of the hormone TSH (thyroid stimulating hormone) [10,11].

Between the peaks of incidence of the disease, two are characteristic, the childhood and the adolescence. In childhood, will be detected to investigate delays in growth or in neuropsychomotor development, caused by hypothyroidism common to pathology. Puberty will be diagnosed as a result of symptomatic manifestations related to the obstruction of the oropharynx, studying with dysphagia, dysphonia, dyspnea, bleeding, pain, feeling of fullness in throat or obstructive sleep apnea. In the newborn, obstructive symptoms occur, there is stridor. In cases of goiter, the main concern is a potential serious obstruction of the upper airways [10,11].

Other situations also correlated to signs of thyroid hypertrophy, which strut asymptomatic, are the States of metabolic stress, such as pregnancy, trauma, surgery, infections, menstruation and menopause [6]. This explains why the lingual thyroid have a higher prevalence in women. Ectopic gland in most cases will provide minimally adequate hormone levels throughout childhood, without causing cretinism [7]. However, with time, will lose the production capacity, being disabled to respond to situations of increased metabolic demands, and hypothyroidism [12].

The initial physical evaluation language includes the thyroid full clinical examination of the head and neck, with special

attention to the oropharynx as lingual thyroid often arises as a formation of submucosal aspect in sessile language basis, median, painless, soft, smooth or irregular surface, highly vascularized and red colored color [11].

The rigid and/or flexible endoscopy allows photo-documentation of the size and position of the lesion, allowing the observation of the larynx and the disposal of other hypotheses, such as cysts of vallecula, hemangiomas and hypertrophy of lingual tonsil. The cervical palpation is essential to search for the presence of thyroid cervical tissue, are not present in general, as has been seen previously. Thus, the clinical diagnosis with the preview directly or with pasta on base red colored laryngoscopy [11-13].

Because of its great food, even trauma vascularization can ulcerate the ectopic tissue, determining copious bleeding. These will offer serious danger to life, by the risk of pulmonary aspiration for the airways [8].

To evaluate the diagnostic criteria, besides the history and physical examination, some additional tests will assist the differential diagnosis, being the study of essential image for final confirmation. The most relevant in this regard will be the mapping with radioactive iodine and technetium99 in bone scans, both showing result of marking the level of the mouth. The first is the most sensitive and specific test for the detection of the existence, size, distribution and activity of ectopic tiroideu fabric and the presence or absence of thyroid gland in its usual location. The second is less specific, but with better cost benefit and is also able to provide important information [8,14].

As in general there will be the absence of tissue in the cervical region, these will be essential exams preoperatively. Avoiding, whenever possible, the total resection of the sole producer of tissue thyroid hormones, what would the patient do hormone replacement for the rest of his life [8,15].

Computed tomography and magnetic resonance imaging are complementary diagnostic methods useful in determining the size of the gland and in surgical planning¹⁶. Sagittal reconstructions will be important in the case of large volume, assessing the depth within the language and air permeability [13].

The ultrasound would differentiate cystic changes and have low cost, but still has a bad role set forth in this pathology, no studies to support its use. The FNA (fine needle aspiration) can also confirm the diagnosis and preoperative examination is the only that can differentiate benign from malignant lesions [16,17]. However, we cannot rule out malignancy in the absence of malignancy puncturing. The definitive diagnosis often is only possible with the ectopic tissue excision [18-20]. In addition to these reasons, the biopsy is also indicated for many authors the high risks of hemorrhage, infection and acute thyrotoxicosis observed puncture [8,13].

Thyroid function tests are also essential in the initial investigation, due to the high line with hypothyroidism, because the hormone produced by the topical is normal, but often insufficient. They do often increased TSH values and

thyroglobulin, and neighboring values or reduced by T3 (triiodothyronine) and T4 (thyroxine) [13-15].

Treatment

There is no consensus in the literature about the best therapeutic strategy, given the rarity of the disease and to the limited number of cases described. Treatment depends on the size, the presence or absence of symptoms and complications like ulceration, hemorrhage, malignancy or obstruction of the upper airway. The fact of ectopic thyroid or may not be the only functional thyroid tissue must also be valued to determine the therapeutic approach. The main objective will be to relieve the obstruction of upper airways without putting the patient's life at risk or leave you with scars mutilating [2,7,21].

Asymptomatic cases can be monitored with serial exams or receive hormonal therapy with levothyroxine aimed at treatment of subclinical hypothyroidism. In addition to treating hormone levels of hypothyroidism, exogenous hormones could cause a suppressive action in gland, reducing its size or preventing a rear hypertrophy. The goal of this treatment would be to suppress the production of TSH, removing the incentive to increase the volume of ectopic gland. This will also be the initial approach of symptomatic patients [2-4,15].

However, such treatment does not have a good success rate. The size reduction speed is very slow, not to expect significant decreases in volume. Many patients gets only the laboratory control, without modification of significant size. Still it is important to note that this therapy can induce acute thyroiditis and sialadenite [4,7,9,16].

Patients treated with hormone suppression should do clinical and laboratory control every 3 months. This time interval can be shortened in predictable situations of increased metabolic demand, such as puberty or pregnancy [7,9,17].

In patients without clinical conditions to be undergoing surgery, those who refused surgical treatment or those in which anatomical difficulties has contraindication the procedure, one can consider treatment with radioactive iodine 131. However, the reduction of thyroid tissue is not consistent, the behavior of the fabric is still unpredictable, and there are descriptions of ectopic tissue that may need higher doses of radioactivity to the ablation. In these cases there is fear to induce radio-induced tumors. So, is completely contraindicated in women of childbearing age and children [2,4,18].

As long as you do not get an adequate response to clinical treatment, the method of surgical thyroidectomy is so preferred. Other indications for surgery are on the basis of urgency: dysphonia or dysphagia, dysphonia important with decreased nutritional intake, suspicion of malignancy, cystic degeneration, uncontrolled hyperthyroidism and repetitive or severe bleeding. Some authors recommend the total thyroidectomy for the possibility of malignant transformation of the lesion [3,7,19].

However, even more experienced hands, total excision of ectopic thyroid tissue is extremely difficult, regardless of the

approach. Remaining scraps of fabric, this will again hypertrophy if replacement is not appropriate or cannot be performed transplantation of tissue removed [4,9,20].

The ectopic tissue autograft tries to avoid hypothyroidism and the replacement with levothyroxine for the rest of his life. If this tissue is grafted or not able to prevent states of hypothyroidism, the fact is hard to predict, which if successful, would prevent the need for replacement therapy. The location for deployment of the fabric is not important in the survival of the same, and the abdominal rectus muscle used most commonly for easy access. It is expected some degree of hypothyroidism first four months postoperatively, but should not be given hormone therapy in order to prevent the failure of the graft [3,4,9,21].

Ensure preoperative airway is crucial, especially in cases of significant obstruction, being achieved through nasotracheal intubation or tracheostomy. The greater the mass, or the laryngeal edema by excessive handling, more prudent will be conducting temporary tracheostomy or, alternatively, keep the nasotracheal intubation for 24 hours after the surgery [3,4,7].

Several surgical approaches have been described, including the trans-hioideas pathways, supra-hioideas, and cervical faringotomy. However, these procedures are largely noninvasive and have high morbidity, leaving large numbers of functional and aesthetic consequences, in addition to the time of hospitalization and recovery period [3,4,7].

Alternatively, a transoral approach offers the advantage of not leaving scars. Most of the cases described in the literature was excised by transoral. The challenge, however, is the difficulty of intraoperative visualization with the naked eye or microscope ectopic thyroid resection of the lesion with a laser and standard surgical instruments. The lack of support of the long surgical instruments and the straight line generated by the laser are the main factors cited to justify this complexity. Another great aggravation of the technique is the control of the intra operative bleeding, and may put the patient's life at risk [2,7,9,16].

Thus, various modifications of the technique were introduced, with the aim of improving exposure, such as incision and opening of the language and of the chin and mandibulotomy. However, some of the reported complications were extensive necrosis of the language, pharyngeal-cutaneous fistula formation and cervical tissue contamination with saliva. Are overly aggressive approaches to the treatment of an injury initially benign [2,3,4,20].

The emergence of transoral robotic-assisted surgery allowed a minimally invasive approach with a larger view of the surgical field and three-dimensional. The endoscopic arm that allows this view base language properly. This system allows the surgeon to employ bi-manual technique, controlling both arms instrument, besides enabling the inclusion of a better Wizard and handling tissue retraction. The greater dexterity and range of motion promoted by robotic equipment allow multiplanar sections within a confined space, due to the flexibility of the neck. These advantages provide a shorter operating time and blood loss, resulting in a surgery more easy, fast and safe for

the patient, including the largest, stretching after the vallecula [17-20].

Conclusion

Lingual Thyroid is a rare congenital anomaly usually asymptomatic, however, when too long causes obstructive symptoms, having your recommended treatment. The approach suppressive hormone should be first line but, in cases of bankruptcy, or if the clinical situation so warrants, the total surgical excision is indicated. The studies evaluated showed great preference for transoral approach, being robotic with the best results in safety, causing minimal morbidity, faster recovery and shorter hospital stay. Only a proper diagnosis of cervical masses can lead to a correct therapy, avoiding iatrogenic.

References

1. Sevinç AI, Unek T, Canda AE, Guray M, Kocdor MA, et al. (2010) Papillary carcinoma arising in subhyoid ectopic thyroid gland with no orthotopic thyroid tissue. *Am J Surg* 200: 17-28.
2. Days, Eady H (2002) Cervical ectopic Thyroid side: case report and review of the literature. *Rev Bras Otorrinolaringol* 68: 128-131.
3. Gu T, Jiang B, Wang N, Xia F, Wang L, et al. (2015) New insight into ectopic thyroid glands between the neck and maxillofacial region from a 42-case study. *BMC Endocr Disord* 15-70.
4. Gotlib J, Ianessi A, Santini J, Dassonville O (2015) Ectopic lingual thyroid. *Eur Ann Otorhinolaryngol Head Neck Dis* 13: 148-149.
5. Hazarika P, Siddiqui SA, Pujary K, Shah P, Nayak DR, et al. (1998) Dual ectopic thyroid: a report of two cases. *J Laryngol Otol* 112: 393-395.
6. Williams JD, Sclafani AP, Slupchinskij O, Douge C (1996) Evaluation and management of the lingual thyroid gland. *Ann Otol Rhinol Laryngol* 105: 312-316.
7. Tincani AJ, Martins AS, Del Negro A, Araujo PP, Barretto G (2004) Lingual thyroid causing dysphonia: evaluation and management. Case report. *Sao Paulo Med J* 122: 67-69.
8. Noussios G, Anagnostis P, Goulis DG, Lappas D, Natsis K (2011) Ectopic thyroid tissue: anatomical, clinical, and surgical implications of a rare entity. *Eur J Endocrinol* 165: 375-382.
9. Banks FAC, Mendes WO, Nunes YP (2014) Ectopic thyroid gland in normal topography submental without. *Rev Bras CIR Head Neck* 43: 172-175.
10. Kumar LK, Kurien NM, Jacob MM, Menon PV, Khalam SA (2015) Lingual thyroid. *Ann Maxillofac Surg* 5: 104-107.
11. Sturniolo G, Violi MA, Galletti B, Baldari S, Campenni A, et al. (2016) Differentiated thyroid carcinoma in lingual thyroid. *Endocrine* 51: 189-198.
12. Chen HW, Lai CC (2015) Lingual thyroid. *J Emerg Med* 48: 103-104.
13. Reed JT, Grant CM (2014) Large mass on base of tongue Diagnosis: lingual thyroid. *J Fam Pract* 63: 603-605.
14. Demir H, Berk F, Isgoren S, Erdinçler RO, Ciftci E, et al. (2002) Subhyoid ectopic thyroid with mediolateral location in a developmentally retarded child with hypothyroidism in an iodine deficiency region. *Ann Nucl Med* 16: 483-486.
15. Atiyeh BS, Abdelnour A, Haddad FF, Ahmad H (1995) Lingual thyroid: tongue-splitting incision for transoral excision. *J Laryngol Otol* 109: 520-524.
16. Prisman E, Patsias A, Genden EM (2015) Transoral robotic excision of ectopic lingual thyroid: Case series and literature review. *Head Neck* 37: 88-91.
17. Pellini R, Mercante G, Ruscito P, Cristalli G, Spriano G (2013) Ectopic lingual goiter treated by transoral robotic surgery. *Acta Otorhinolaryngol Ital* 33: 343-346.
18. Teo EH, Toh ST, Tay HN, Han HJ (2013) Transoral robotic resection of lingual thyroid: case report. *J Laryngol Otol* 127: 1034-1037.
19. Byrd JK, Duvvuri U (2013) Current trends in robotic surgery for otolaryngology. *Curr Otorhinolaryngol Rep* 1: 153-157.
20. Park YM, Kim WS, Byeon HK, Lee SY, Kim SH (2013) A novel technique for the resection of the symptomatic lingual thyroid: transoral robotic surgery. *Thyroid* 23: 466-471.
21. Amani Mel A, Benabadji N, Benzian Z, Amani S (2012) Ectopic lingual thyroid. *Indian J Nucl Med* 27: 124-126.