

Paris. France

April 18-19, 2019

World Congress on

Immuno Oncology & **Clinical Pharmacy**

Arch Cancer Res 2019, Volume:7 DOI: 10.21767/2254-6081-C2-024

NECROTIZING ENTEROCOLITIS IN A PRETERM INFANT NEWBORN AND ROLE OF FEEDING: AN UPDATE

Amr I M Hawal

Prime Health Care Group, UAE

Statement of the problem: It's a clinical case presentation of a male preterm infant new born (+31 weeks) who was delivered in our hospital and transferred to our NICU because of prematurity, VLBW and need respiratory support. Baby shortly undergo Necrotizing enterocolitis (NEC) on 5th day of life shortly after start of expressed milk feeding, which was early detected by use of near infrared abdominal spectroscopy (NIRS). Baby was deteriorated clinically in a couple of hours and undergone intestinal perforation with peritonitis. So, the abdominal exploration surgery with intestinal resection and end to end anastomosis was done urgently. Baby was improved gradually. Early feedings was started and gradually increased up to full feedings with use of human fortified milk (HMF) and probiotics, Prebiotics.

Findings: The Study stated the evidence-based feeding strategies guidelines for necrotizing enterocolitis (NEC) among very low birth weight infants and the role of trophic feedings, probiotics, prebiotics and micronutrients in prophylaxis, prevention and management of NEC.

Recommendations: Prematurity is the single greatest risk factor for NEC and avoidance of premature birth is the best way to prevent NEC; The role of feeding in the pathogenesis of NEC is uncertain, but it seems prudent to use breast milk (when available) and advance feedings slowly and cautiously; NEC is one of the leading causes of mortality and the most common reason for emergent GI surgery in new borns; NEC remains a major unsolved medical challenge, for which no specific therapy exists and its pathogenesis remains controversial; A better understanding of the pathophysiology will offer new and innovative therapeutic approaches and future studies should be focused on the roles of the epithelial barrier, innate immunity and microbiota in this disorder; Bioinformatics modelling is a new emerging strategy aimed at understanding the dynamics of various inflammatory markers and their application in early diagnosis and treatment.

amr106@gmail.com