

ATHEROSCLEROTIC CHANGES IN PATIENTS WITH OBSTRUCTIVE SLEEP APNEA

**Victor Manolov¹, Georgiev O², Vasilev V³, Pencheva Genova V²,
Grozdanova R⁴, Traykov L⁵, Petrova I⁵, Tzatchev K¹, Hadjideko-
va S⁶, Karadjova M⁵, Voleva S⁷, Nikolova M⁴, Angov G⁵, Petrova
Ivanova I⁵, Kunchev T⁵, Ovcharov D⁵, Gramatikova Z⁸**

¹Dept. of Clinical Laboratory and Clinical Immunology, Medical University, Bulgaria

²Dept. of Propedeutics of Internal Diseases, Medical University Sofia

³Clinical laboratory and clinical pharmacology, University Aleksandrovska hospital Sofia, Bulgaria

⁴NCIP, Bulgaria

⁵Department of Neurology, Medical University Sofia

⁶Dept. of Medical Genetics, Medical University Sofia

⁷ NCIPD, Sofia

⁸RED Laboratories N V/SA, Belgium

O bstructive sleep apnea syndrome (OSA) is defined as a combination of symptoms as a result of intermittent, recurrent constraint and / or complete airway overhead airway overflow (sleep disturbance). During desaturation episodes, the organism is subjected to chronic stress. This leads to reduced nitric oxide secretion, increased release of interleukin-6, tumour necrosis factor-alpha and other pro-inflammatory cytokines. The described pathological cascades are associated with the development of insulin resistance, arterial hypertension, metabolic syndrome, systemic atherosclerosis and increased cardiovascular risk. 47 patients aged 43.1±7.7 with OSA were included in this study. Their results were compared to sex and age matched healthy control. CBC, serum iron, ferritin, hsCRP, hepcidin, homocysteine and vitamin B12 were measured in the included groups. IMT and FMT were used for atherosclerotic changes evaluation. We found increased serum hepcidin levels in OSA patients with IMT and FMD changes (99.8±14.5 µg/L) compared to healthy controls (19.7±1.9 µg/L); P<0.001. IMT and FMD correlates positive in OSA patients with atherosclerotic changes to serum hepcidin concentrations (r=0.811, r=0.847, resp.; P<0.01). Serum hepcidin correlates positively to homocysteine and hsCRP in OSA patients (r=0.788, r=0.797, resp.; P<0.005). Brain-vascular disease risk factors are connected to obstructive sleep apnea syndrome. Disregulation of iron homeostasis is one of the main risk atherogenesis factors. Early hepcidin quantification might predict an atherosclerosis occurrence in OSA patients, which might be very important for better clinical diagnosis and practice.

vichedoc4@abv.bg